arXiv:1304.4894v1 [math.CA] 15 Apr 2013

SOME HADAMARD LIKE INEQUALITIES VIA CONVEX AND s-CONVEX FUNCTIONS AND THEIR APPLICATIONS FOR SPECIAL MEANS

MEVLÜT TUNÇ

Abstract. In this study, the author establish some inequalities of Hadamard like based on convex and s-convexity in the second sense. Some applications to special means of positive real numbers are also given.

1. Preliminaries

1.1. Definitions.

Definition 1. [15] *A function* $f : I \to \mathbb{R}$ *is said to be convex on* I *if inequality*

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) \tag{1.1}$$

holds for all $x, y \in I$ *and* $t \in [0,1]$. *We say that* f *is concave if* $(-f)$ *is convex.*

Geometrically, this means that if P, Q and R are three distinct points on the graph of f with Q between P and R, then Q is on or below the chord PR.

Definition 2. [11] *Let* $s \in (0,1]$. *A function* $f : (0,\infty] \to [0,\infty]$ *is said to be* s*-convex in the second sense if*

$$f(tx + (1-t)y) \leq t^s f(x) + (1-t)^s f(y), \tag{1.2}$$

for all $x, y \in (0,b]$ *and* $t \in [0,1]$. *This class of* s*-convex functions is usually denoted by* K_s^2.

Certainly, s-convexity means just ordinary convexity when $s = 1$.

1.2. Theorems.

Theorem 1. ***The Hermite-Hadamard inequality:*** *Let* $f : I \subseteq \mathbb{R} \to \mathbb{R}$ *be a convex function and* $u, v \in I$ *with* $u < v$. *The following double inequality:*

$$f\left(\frac{u+v}{2}\right) \leq \frac{1}{v-u}\int_u^v f(x)\,dx \leq \frac{f(u)+f(v)}{2} \tag{1.3}$$

is known in the literature as Hadamard's inequality (or Hermite-Hadamard inequality) for convex functions. If f *is a positive concave function, then the inequality is reversed.*

2000 *Mathematics Subject Classification.* Primary 26A15, 26D10; Secondary 26D07, 26D15.
Key words and phrases. Hadamard' inequality, convexity, s-convexity.
This paper is in final form and no version of it will be submitted for publication elsewhere.

Theorem 2. [8] *Suppose that* $f:[0,\infty)\rightarrow[0,\infty)$ *is an* $s-$*convex function in the second sense, where* $s\in(0,1]$, *and let* $a,b\in[0,\infty)$, $a<b$. *If* $f\in L_1([0,1])$, *then the following inequalities hold:*

$$2^{s-1}f\left(\frac{u+v}{2}\right)\leq\frac{1}{v-u}\int_u^v f(x)\,dx\leq\frac{f(u)+f(v)}{s+1}. \tag{1.4}$$

The constant $k=\frac{1}{s+1}$ *is the best possible in the second inequality in (1.4). The above inequalities are sharp. If* f *is an s-concave function in the second sense, then the inequality is reversed.*

For recent results and generalizations concerning Hadamard's inequality and concepts of convexity and s-convexity see [1]-[20] and the references therein.

Throughout this paper we will use the following notations and conventions. Let $J=[0,\infty)\subset\mathbb{R}=(-\infty,+\infty)$, and $u,v\in J$ with $0<u<v$ and $f'\in L[u,v]$ and

$$\begin{aligned} A(u,v) &= \frac{u+v}{2},\ G(u,v)=\sqrt{uv},\ I(u,v)=\frac{1}{e}\left(\frac{b^b}{a^a}\right)^{\frac{1}{b-a}}, \\ L_p(u,v) &= \left(\frac{v^{p+1}-u^{p+1}}{(p+1)(v-u)}\right)^{1/p},\ u\neq v,\ p\in\mathbb{R},\ p\neq-1,0 \end{aligned}$$

be the arithmetic mean, geometric mean, identric mean, generalized logarithmic mean for $u,v>0$ respectively.

2. Some new Hadamard like inequalities

In order to establish our main results, we first establish the following lemma.

Lemma 1. *Let* $f:J\rightarrow\mathbb{R}$ *be a differentiable function on* J°. *If* $f'\in L[u,v]$, *then*

$$\begin{aligned} &\frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu \\ =\ &\frac{(v-x)^2}{(v-u)^2}\int_0^1(tu+(1-t)v)f'(tx+(1-t)v)\,dt \\ &+\frac{(x-u)^2}{(v-u)^2}\int_0^1(tv+(1-t)u)f'(tx+(1-t)u)\,dt \end{aligned}$$

for each $t\in[0,1]$ *and* $x\in[u,v]$.

Proof. Integrating by parts, we get

$$\begin{aligned}
& \frac{(v-x)^2}{(v-u)^2}\int_0^1 (tu+(1-t)v)\, f'(tx+(1-t)v)\,dt \\
& +\frac{(x-u)^2}{(v-u)^2}\int_0^1 (tv+(1-t)u)\, f'(tx+(1-t)u)\,dt \\
= & \frac{(v-x)^2}{(v-u)^2}\left[(tu+(1-t)v)\left.\frac{f(tx+(1-t)v)}{x-v}\right|_0^1 - (u-v)\int_0^1 \frac{f(tx+(1-t)v)}{x-v}dt\right] \\
& +\frac{(x-u)^2}{(v-u)^2}\left[(tv+(1-t)u)\left.\frac{f(tx+(1-t)u)}{x-u}\right|_0^1 - (v-u)\int_0^1 \frac{f(tx+(1-t)u)}{x-u}dt\right] \\
= & \frac{(v-x)^2}{(v-u)^2}\left[\frac{uf(x)-vf(v)}{x-v}-\frac{v-u}{(x-v)^2}\int_x^v f(\mu)\,d\mu\right] \\
& +\frac{(x-a)^2}{(v-u)^2}\left[\frac{vf(x)-uf(u)}{x-u}-\frac{v-u}{(x-u)^2}\int_u^x f(\mu)\,d\mu\right] \\
= & \frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu.
\end{aligned}$$

□

Theorem 3. *Let $f: J \to \mathbb{R}$ be a differentiable function on J°. If $|f'|$ is convex on $[u,v]$, then*

$$\begin{aligned}
& \left|\frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq & \frac{(v-x)^2}{6(v-u)^2}\left[(2u+v)\left|f'(x)\right|+(u+2v)\left|f'(v)\right|\right]+\frac{(x-u)^2}{6(v-u)^2}\left[(u+2v)\left|f'(x)\right|+(2u+v)\left|f'(u)\right|\right]
\end{aligned}$$

for each $x \in [u,v]$.

Proof. Using Lemma 1 and from properties of modulus, and since $|f'|$ is convex on $[u,v]$, then we obtain

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\int_0^1 (tu+(1-t)v)\left|f'(tx+(1-t)v)\right|dt \\
&+\frac{(x-u)^2}{(v-u)^2}\int_0^1 (tv+(1-t)u)\left|f'(tx+(1-t)u)\right|dt \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\int_0^1 (tu+(1-t)v)\left(t\left|f'(x)\right|+(1-t)\left|f'(v)\right|\right)dt \\
&+\frac{(x-u)^2}{(v-u)^2}\int_0^1 (tv+(1-t)u)\left(t\left|f'(x)\right|+(1-t)\left|f'(u)\right|\right)dt \\
=\ & \frac{(v-x)^2}{6(v-u)^2}\left[(2u+v)\left|f'(x)\right|+(u+2v)\left|f'(v)\right|\right] \\
&+\frac{(x-u)^2}{6(v-u)^2}\left[(u+2v)\left|f'(x)\right|+(2u+v)\left|f'(u)\right|\right].
\end{aligned}$$

The proof is completed. □

Theorem 4. *Let $f: J\to\mathbb{R}$ be a differentiable function on J°. If $|f'|$ is s-convex on $[u,v]$ for some fixed $s\in(0,1]$, then*

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\left[\frac{((s+1)u+v)\left|f'(x)\right|+(u+(s+1)v)\left|f'(v)\right|}{(s+1)(s+2)}\right] \\
&+\frac{(x-u)^2}{(v-u)^2}\left[\frac{(u+(s+1)v)\left|f'(x)\right|+((s+1)v+v)\left|f'(u)\right|}{(s+1)(s+2)}\right]
\end{aligned}$$

for each $x\in[u,v]$.

Proof. The proof of Theorem 4 is similar to Theorem 3. □

Remark 1. *In Theorem 3, if we take $s=1$, then Theorem 4 reduces to Theorem 3.*

Corollary 1. *In Theorem 4, if we choose $x=\frac{u+v}{2}$, we get*

$$\begin{aligned}
&\left|\frac{vf(v)-uf(u)}{2(v-u)}+\frac{1}{2}f\left(\frac{u+v}{2}\right)-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq\ & \frac{1}{4}\left[\frac{((s+1)u+v)\left|f'\left(\frac{u+v}{2}\right)\right|+(u+(s+1)v)\left|f'(v)\right|}{(s+1)(s+2)}\right] \\
&+\frac{1}{4}\left[\frac{(u+(s+1)v)\left|f'\left(\frac{u+v}{2}\right)\right|+((s+1)u+v)\left|f'(u)\right|}{(s+1)(s+2)}\right]
\end{aligned}$$

Proposition 1. *Let* $u, v \in J^{\circ},\ 0<u<v$ *and* $s \in (0,1]$, *then*

$$\begin{aligned}
&\left|\frac{s+1}{2}L_s^s(u,v)+\frac{1}{2}A^s(u,v)-L_s^s(u,v)\right| \\
= &\left|(s-1)L_s^s(u,v)+A^s(u,v)\right| \qquad (2.1)\\
\leq &\frac{s}{s+1}A(u,v)+\frac{s}{s+2}A(u^s,v^s)+\frac{sG^2(u,v)}{(s+1)(s+2)}A\left(u^{s-2},v^{s-2}\right)
\end{aligned}$$

Proof. The proof follows from Corollary 1 applied to the s-convex function $f(x)=x^s$. Equality in (2.1) holds if and only if $s=1$. ☐

Corollary 2. *In Corollary 1, if we take* $|f'| \leq M$, *we get*

$$\left|\frac{vf(v)-uf(u)}{2(v-u)}+\frac{1}{2}f\left(\frac{u+v}{2}\right)-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \leq M\frac{(u+v)}{2(s+1)}$$

Theorem 5. *Let* $f: J \to \mathbb{R}$ *be a differentiable function on* J°. *If* $|f'|^q$ *is convex on* $[u,v]$ *and* $q>1$ *with* $1/p+1/q=1$, *then*

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq &\frac{(v-x)^2}{(v-u)^2}L_p(u,v)\left(\frac{|f'(x)|^q+|f'(v)|^q}{2}\right)^{\frac{1}{q}}+\frac{(x-u)^2}{(v-u)^2}L_p(u,v)\left(\frac{|f'(x)|^q+|f'(u)|^q}{2}\right)^{\frac{1}{q}}.
\end{aligned}$$

for each $x \in [u,v]$.

Proof. Using Lemma 1 and using the well-known Hölder's inequality and since $|f'|^q$ is convex on $[u,v]$, we establish

(2.2)

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq &\frac{(v-x)^2}{(v-u)^2}\int_0^1 (tu+(1-t)v)\left|f'(tx+(1-t)v)\right|dt \\
&+\frac{(x-u)^2}{(v-u)^2}\int_0^1 (tv+(1-t)u)\left|f'(tx+(1-t)u)\right|dt \\
\leq &\frac{(v-x)^2}{(v-u)^2}\left(\int_0^1 (tu+(1-t)v)^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 \left|f'(tx+(1-t)v)\right|^q dt\right)^{\frac{1}{q}} \\
&+\frac{(x-u)^2}{(v-u)^2}\left(\int_0^1 (tv+(1-t)u)^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 \left|f'(tx+(1-t)u)\right|^q dt\right)^{\frac{1}{q}}.
\end{aligned}$$

Since $|f'|^q$ is convex on $[u,v]$, by Hadamard's inequality, we have

$$\int_0^1 \left|f'(tx+(1-t)v)\right|^q dt \leq \frac{|f'(x)|^q+|f'(v)|^q}{2} \qquad (2.3)$$

$$\int_0^1 \left|f'(tx+(1-t)u)\right|^q dt \leq \frac{|f'(x)|^q+|f'(u)|^q}{2} \qquad (2.4)$$

It can be easily seen that

$$\int_0^1 (tu+(1-t)\,v)^p\,dt=\int_0^1 (tv+(1-t)\,u)^p\,dt=\frac{v^{p+1}-u^{p+1}}{(v-u)\,(p+1)}=L_p^p\,(u,v) \tag{2.5}$$

If expressions (2.3)-(2.5) are written in (2.2), we obtain

$$\begin{aligned}&\left|\frac{(v-x)\,(vf\,(v)-uf\,(x))+(x-u)\,(vf\,(x)-uf\,(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f\,(\mu)\,d\mu\right|\\ \leq\ &\frac{(v-x)^2}{(v-u)^2}L_p\,(u,v)\left(\frac{|f'\,(x)|^q+|f'\,(v)|^q}{2}\right)^{\frac{1}{q}}+\frac{(x-u)^2}{(v-u)^2}L_p\,(u,v)\left(\frac{|f'\,(x)|^q+|f'\,(u)|^q}{2}\right)^{\frac{1}{q}}.\end{aligned}$$

The proof is completed. □

Theorem 6. *Let $f:J\to\mathbb{R}$ be a differentiable function on J°. If $|f'|^q$ is s-convex on $[u,v]$ for some fixed $s\in(0,1]$ and $q>1$ with $1/p+1/q=1$, then*

$$\begin{aligned}&\left|\frac{(v-x)\,(vf\,(v)-uf\,(x))+(x-u)\,(vf\,(x)-uf\,(u))}{(v-u)^2}-\frac{1}{v-u}\int_u^v f\,(\mu)\,d\mu\right|\\ \leq\ &\frac{(v-x)^2}{(v-u)^2}L_p\,(u,v)\left(\frac{|f'\,(x)|^q+|f'\,(v)|^q}{s+1}\right)^{\frac{1}{q}}+\frac{(x-u)^2}{(v-u)^2}L_p\,(u,v)\left(\frac{|f'\,(x)|^q+|f'\,(u)|^q}{s+1}\right)^{\frac{1}{q}}.\end{aligned}$$

for each $x\in[u,v]$.

Proof. The proof of Theorem 6 is similar to Theorem 5. □

Corollary 3. *In Theorem 6,*

i) if we take $x=u$ or $x=v$, we get

$$\left|\frac{vf\,(v)-uf\,(u)}{v-u}-\frac{1}{v-u}\int_u^v f\,(\mu)\,d\mu\right|\leq L_p\,(u,v)\left(\frac{|f'\,(u)|^q+|f'\,(v)|^q}{s+1}\right)^{\frac{1}{q}}. \tag{2.6}$$

ii) if we take $x=\frac{u+v}{2}$ and since $\left(\frac{1}{s+1}\right)^{\frac{1}{q}}\leq 1,\ s\in(0,1]$, we get

$$\left|\frac{vf\,(v)-uf\,(u)}{2\,(v-u)}+\frac{1}{2}f\left(\frac{u+v}{2}\right)-\frac{1}{v-u}\int_u^v f\,(\mu)\,d\mu\right|$$

$$\leq\frac{L_p\,(u,v)}{4}\left[\left(\frac{\left|f'\left(\frac{u+v}{2}\right)\right|^q+|f'\,(v)|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{u+v}{2}\right)\right|^q+|f'\,(u)|^q}{s+1}\right)^{\frac{1}{q}}\right] \tag{2.7}$$

$$\leq\frac{L_p\,(u,v)}{4}\left[\left(\left|f'\left(\frac{u+v}{2}\right)\right|^q+|f'\,(v)|^q\right)^{\frac{1}{q}}+\left(\left|f'\left(\frac{u+v}{2}\right)\right|^q+|f'\,(u)|^q\right)^{\frac{1}{q}}\right] \tag{2.8}$$

Proposition 2. *Let $u,v\in J^{\circ},\ 0<u<v$ and $s\in(0,1]$, then*

$$\begin{aligned}&\left|(s-1)\,L_s^s\,(u,v)+A^s\,(u,v)\right|\\ \leq\ &\frac{L_p\,(u,v)}{2}\left(\frac{s}{s+1}\right)^{\frac{1}{q}}\left[\left(\left(\frac{u+v}{2}\right)^{(s-1)q}+v^{(s-1)q}\right)^{\frac{1}{q}}+\left(\left(\frac{u+v}{2}\right)^{(s-1)q}+u^{(s-1)q}\right)^{\frac{1}{q}}\right]\end{aligned}$$

Proof. The proof follows from (2.7) applied to the s-convex function $f\,(x)=x^s$. Equality in (2.1) holds if and only if $s=1$ and $u=v$. □

Theorem 7. *Let $f : J \to \mathbb{R}$ be a differentiable function on J°. If $|f'|^q$ is s-convex on $[u, v]$ for some fixed $s \in (0, 1]$ and $q \geq 1$, then*

$$\begin{aligned}
&\left| \frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu \right| \\
\leq\ & \frac{(v-x)^2}{(v-u)^2} A^{1-\frac{1}{q}}(u,v) \left(\frac{1}{(s+1)(s+2)}\right)^{\frac{1}{q}} \left(((s+1)u+v)|f'(x)|^q + ((s+1)v+u)|f'(v)|^q\,dt\right)^{\frac{1}{q}} \\
&+ \frac{(x-u)^2}{(v-u)^2} A^{1-\frac{1}{q}}(u,v) \left(\frac{1}{(s+1)(s+2)}\right)^{\frac{1}{q}} \left(((s+1)v+u)|f'(x)|^q + ((s+1)u+v)|f'(u)|^q\,dt\right)^{\frac{1}{q}}
\end{aligned}$$

for each $x \in [u, v]$.

Proof. Using Lemma 1 and the well-known power mean inequality and since $|f'|^q$ is s-convex on $[u, v]$, we establish

$$\begin{aligned}
&\left| \frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu \right| \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\int_0^1 (tu+(1-t)v)\,|f'(tx+(1-t)v)|\,dt + \frac{(x-u)^2}{(v-u)^2}\int_0^1 (tv+(1-t)u)\,|f'(tx+(1-t)u)|\,dt \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\left(\int_0^1 (tu+(1-t)v)\,dt\right)^{1-\frac{1}{q}} \left(\int_0^1 (tu+(1-t)v)\,|f'(tx+(1-t)v)|^q\,dt\right)^{\frac{1}{q}} \\
&+ \frac{(x-u)^2}{(v-u)^2}\left(\int_0^1 (tv+(1-t)u)\,dt\right)^{1-\frac{1}{q}} \left(\int_0^1 (tv+(1-t)u)\,|f'(tx+(1-t)u)|^q\,dt\right)^{\frac{1}{q}} \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\left(\int_0^1 (tu+(1-t)v)\,dt\right)^{1-\frac{1}{q}} \left(\int_0^1 (tu+(1-t)v)\left(t^s|f'(x)|^q+(1-t)^s|f'(v)|^q\right)dt\right)^{\frac{1}{q}} \\
&+ \frac{(x-u)^2}{(v-u)^2}\left(\int_0^1 (tv+(1-t)u)\,dt\right)^{1-\frac{1}{q}} \left(\int_0^1 (tv+(1-t)u)\left(t^s|f'(x)|^q+(1-t)^s|f'(u)|^q\right)dt\right)^{\frac{1}{q}} \\
=\ & \frac{(v-x)^2}{(v-u)^2}\left(\frac{u+v}{2}\right)^{1-\frac{1}{q}} \left(\frac{(s+1)u+v}{(s+1)(s+2)}|f'(x)|^q + \frac{(s+1)v+u}{(s+1)(s+2)}|f'(v)|^q\,dt\right)^{\frac{1}{q}} \\
&+ \frac{(x-u)^2}{(v-u)^2}\left(\frac{u+v}{2}\right)^{1-\frac{1}{q}} \left(\frac{(s+1)v+u}{(s+1)(s+2)}|f'(x)|^q + \frac{(s+1)u+v}{(s+1)(s+2)}|f'(u)|^q\,dt\right)^{\frac{1}{q}}
\end{aligned}$$

The proof is completed. □

Theorem 8. *Let $f : J \to \mathbb{R}$ be a differentiable function on J°. If $|f'|^q$ is convex on $[u, v]$ and $q \geq 1$, then*

$$\begin{aligned}
&\left| \frac{(v-x)(vf(v)-uf(x))+(x-u)(vf(x)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu \right| \\
\leq\;& \frac{(v-x)^2}{(v-u)^2} A^{1-\frac{1}{q}}(u,v)\left(\frac{1}{6}\right)^{\frac{1}{q}} \left((2u+v)\left|f'(x)\right|^q + (2v+u)\left|f'(v)\right|^q dt\right)^{\frac{1}{q}} \\
&+\frac{(x-u)^2}{(v-u)^2} A^{1-\frac{1}{q}}(u,v)\left(\frac{1}{6}\right)^{\frac{1}{q}} \left((2v+u)\left|f'(x)\right|^q + (2u+v)\left|f'(u)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

for each $x \in [u, v]$.

Proof. In Theorem 7, if we take $s = 1$, then the assertion is proved. ☐

Corollary 4. *In Theorem 7,*

i) if we take $x = u$ or $x = v$, we get

$$\begin{aligned}
&\left| \frac{(v-u)(vf(v)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu \right| \\
\leq\;& A^{1-\frac{1}{q}}(u,v)\left(\frac{1}{(s+1)(s+2)}\right)^{\frac{1}{q}} \left(((s+1)u+v)\left|f'(u)\right|^q + ((s+1)v+u)\left|f'(v)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

ii) if we choose $x = u$ or $x = v$ and $s = q = 1$, we get

$$\left| \frac{(v-u)(vf(v)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu \right| \leq \frac{1}{6}\left((2u+v)\left|f'(u)\right| + (2v+u)\left|f'(v)\right| dt\right)$$

iii) if we take $x = \frac{u+v}{2}$ and since $\left(\frac{1}{(s+1)(s+2)}\right)^{\frac{1}{q}} \leq 1$, $s \in (0, 1]$, we get

$$\begin{aligned}
&\left| \frac{vf(v)-uf(u)}{2(v-u)} + \frac{1}{2}f\left(\frac{u+v}{2}\right) - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu \right| \\
&\leq \frac{1}{4}A^{1-\frac{1}{q}}(u,v)\left(\frac{1}{(s+1)(s+2)}\right)^{\frac{1}{q}} \left(((s+1)u+v)\left|f'\left(\frac{u+v}{2}\right)\right|^q + ((s+1)v+u)\left|f'(v)\right|^q dt\right)^{\frac{1}{q}} \\
&\quad+\frac{1}{4}A^{1-\frac{1}{q}}(u,v)\left(\frac{1}{(s+1)(s+2)}\right)^{\frac{1}{q}} \left(((s+1)v+u)\left|f'\left(\frac{u+v}{2}\right)\right|^q + ((s+1)u+v)\left|f'(u)\right|^q dt\right)^{\frac{1}{q}} \qquad (2.9) \\
&\leq \frac{1}{4}A^{1-\frac{1}{q}}(u,v)\left(((s+1)u+v)\left|f'\left(\frac{u+v}{2}\right)\right|^q + ((s+1)v+u)\left|f'(v)\right|^q dt\right)^{\frac{1}{q}} \qquad (2.10) \\
&\quad+\frac{1}{4}A^{1-\frac{1}{q}}(u,v)\left(((s+1)v+u)\left|f'\left(\frac{u+v}{2}\right)\right|^q + ((s+1)u+v)\left|f'(u)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

Proposition 3. *Let* $u, v \in J^{\circ},\ 0 < u < v$ *and* $s \in (0,1]$, *then*

$$\begin{aligned}
&\left|(s-1) L_s^s(u,v) + A^s(u,v)\right| \\
\leq\ & \left(\frac{s^q A^{q-1}(u,v)}{(s+1)(s+2)2^q}\right)^{\frac{1}{q}} \left(((s+1)u+v) A^{(s-1)q}(u,v) + ((s+1)v+u) v^{(s-1)q} dt\right)^{\frac{1}{q}} \\
&+\left(\frac{s^q A^{q-1}(u,v)}{(s+1)(s+2)2^q}\right)^{\frac{1}{q}} \left(((s+1)v+u) A^{(s-1)q}(u,v) + ((s+1)u+v) u^{(s-1)q} dt\right)^{\frac{1}{q}}
\end{aligned}$$

Proof. The proof follows from (2.9) applied to the s-convex function $f(x) = x^s$. $\square$

Theorem 9. *Let* $f : J \to \mathbb{R}$ *be a differentiable function on* J°. *If* $|f'|^q$ *is* s*-concave on* $[u,v]$ *for some fixed* $s \in (0,1]$ *and* $q > 1$ *with* $1/p + 1/q = 1$, *then*

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x)) + (x-u)(vf(x)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq\ & \frac{2^{\frac{s-1}{q}} L_p(u,v)}{(v-u)^2}\left[(v-x)^2\left|f'\left(\frac{x+v}{2}\right)\right| + (x-u)^2\left|f'\left(\frac{x+u}{2}\right)\right|\right]
\end{aligned}$$

for each $x \in [u,v]$.

Proof. Using Lemma 1 and using the Hölder inequality and since $|f'|^q$ is s-concave on $[u,v]$ and using the inequality (1.4), we establish

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x)) + (x-u)(vf(x)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\int_0^1 (tu+(1-t)v)\left|f'(tx+(1-t)v)\right| dt + \frac{(x-u)^2}{(v-u)^2}\int_0^1 (tv+(1-t)u)\left|f'(tx+(1-t)u)\right| dt \\
\leq\ & \frac{(v-x)^2}{(v-u)^2}\left(\int_0^1 (tu+(1-t)v)^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 \left|f'(tx+(1-t)v)\right|^q dt\right)^{\frac{1}{q}} \\
&+\frac{(x-u)^2}{(v-u)^2}\left(\int_0^1 (tv+(1-t)u)^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 \left|f'(tx+(1-t)u)\right|^q dt\right)^{\frac{1}{q}} \\
\leq\ & \frac{(v-x)^2}{(v-u)^2} L_p(u,v)\left(2^{s-1}\left|f'\left(\frac{x+v}{2}\right)\right|^q\right)^{\frac{1}{q}} + \frac{(x-u)^2}{(v-u)^2} L_p(u,v)\left(2^{s-1}\left|f'\left(\frac{x+u}{2}\right)\right|^q\right)^{\frac{1}{q}}
\end{aligned}$$

The proof is completed. $\square$

Theorem 10. *Let* $f : J \to \mathbb{R}$ *be a differentiable function on* J°. *If* $|f'|^q$ *is concave on* $[u,v]$ *and* $q > 1$ *with* $1/p + 1/q = 1$, *then*

$$\begin{aligned}
&\left|\frac{(v-x)(vf(v)-uf(x)) + (x-u)(vf(x)-uf(u))}{(v-u)^2} - \frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \\
\leq\ & \frac{L_p(u,v)}{(v-u)^2}\left[(v-x)^2\left|f'\left(\frac{x+v}{2}\right)\right| + (x-u)^2\left|f'\left(\frac{x+u}{2}\right)\right|\right]
\end{aligned}$$

for each $x \in [u,v]$.

Proof. Using Lemma 1 and using the Hölder inequality and since $|f'|^q$ is concave on $[u,v]$ and using Hadamard's inequality for concave functions, we complete the proof. Or, in Theorem 9, if we take $s=1$, then the assertion is also proved. □

Corollary 5. *In Theorem 9,*

i) if we take $x=u$ or $x=v$, we get

$$\left|\frac{(v-u)\left(vf(v)-uf(u)\right)}{(v-u)^2}-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \leq 2^{\frac{s-1}{q}} L_p(u,v)\left|f'\left(\frac{u+v}{2}\right)\right|$$

ii) if we take $x=\frac{u+v}{2}$, we get

$$\left|\frac{vf(v)-uf(u)}{2(v-u)}+\frac{1}{2}f\left(\frac{u+v}{2}\right)-\frac{1}{v-u}\int_u^v f(\mu)\,d\mu\right| \leq \frac{2^{\frac{s-1}{q}}L_p(u,v)}{4}\left[\left|f'\left(\frac{u+3v}{4}\right)\right|+\left|f'\left(\frac{v+3u}{4}\right)\right|\right] \tag{2.11}$$

Proposition 4. *Let $u,v\in J^{\circ}$, $0<u<v$ and $s=1$, then*

$$\left|\frac{v\sin v-u\sin u+2\cos v-2\cos u}{v-u}+\sin\left(A(u,v)\right)\right| \leq \frac{L_p(u,v)}{2}\left[\left|\cos\left(\frac{u+3v}{4}\right)\right|+\left|\cos\left(\frac{v+3u}{4}\right)\right|\right]$$

Proof. The proof follows from (2.11) applied to the concave function $f:[0,\pi]\to[0,1]$, $f(x)=\sin x$. □

References

[1] M. Alomari, M. Darus, U.S. Kırmacı, *Some Inequalities of Hermite-Hadamard type for s-convex Functions*, *Acta Math. Sci.* 31B(4), (2011)1643-1652.

[2] M. Avci, H. Kavurmaci, M. E. Özdemir, *New inequalities of Hermite-Hadamard type via s-convex functions in the second sense with applications*, Appl. Math. Comp. 217 (2011) 5171-5176.

[3] W. W. Breckner, *Stetigkeitsaussagen f¨ur eine Klasse verallgemeinerter konvexer funktionen in topologischen linearen Raumen*, Pupl. Inst. Math., 23, 13-20, (1978).

[4] W. W. Breckner, *Continuity of generalized convex and generalized concave set-valued functions*, Rev Anal. Num´er. Thkor. Approx., 22, 39–51, (1993).

[5] P. Burai, A. Házy, and T. Juhász, *Bernstein–Doetsch type results for s-convex functions*, Publ. Math. Debrecen 75 (2009), no. 1-2, 23-31.

[6] P. Burai, A. Házy, and T. Juhász, *On approximately Breckner s-convex functions*, Control Cybernet. 40 (2011), no. 1, 91-99.

[7] S.S. Dragomir, R.P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, *Appl. Math. Lett.* 11 (1998) 91-95.

[8] S.S. Dragomir, S. Fitzpatrick, *The Hadamard's inequality for s-convex functions in the second sense*, Demonstration Math., 32 (4) (1999), 687–696.

[9] S. S. Dragomir and C. E. M. Pearce: *Selected topics on Hermite-Hadamard inequalities and applications, RGMIA monographs, Victoria University*, 2000. [Online: http://www.staff.vu.edu.au/RGMIA/monographs/hermite-hadamard.html].

[10] J. Hadamard, *Étude sur les propriétés des fonctions entières en particulier d'une fonction considérée par Riemann*, J. Math. Pures Appl. 58 (1893) 171–215.

[11] H. Hudzik and L. Maligranda, *Some remarks on s-convex functions*, Aequationes Math., 48, 100-111, (1994).

[12] H. Kavurmaci, M. Avci, M E. Özdemir, *New inequalities of hermite-hadamard type for convex functions with applications,* Jour. Ineq. Appl. 2011, 2011:86, doi:10.1186/1029-242X-2011-86.

[13] U.S. Kırmacı, M.E. Özdemir, *On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula*, Appl. Math. Comput. 153 (2004) 361-368.

[14] D.S. Mitrinović, I.B. Lacković, *Hermite and convexity*, Aequationes Math. 28 (1985) 229-232.

[15] D. S. Mitrinović, J. Pečarić, and A.M. Fink, *Classical and new inequalities in analysis*, KluwerAcademic, Dordrecht, 1993.

[16] M.E. Özdemir, *A theorem on mappings with bounded derivatives with applications to quadrature rules and means,* Appl. Math. Comput. 138 (2003) 425-434.

[17] J. E. Pečarić, F. Proschan and Y. L. Tong: *Convex Functions, Partial Orderings, and Statistical Applications, Academic Press Inc.,* 1992.

[18] M.Z. Sarikaya, E. Set, M.E. Özdemir, *On new inequalities of Simpson's type for s-convex functions*, Comp&Math Appl., 60 (2010) 2191-2199.

[19] B.-Y. Xi and F. Qi, *Some Integral Inequalities of Hermite-Hadamard Type for Convex Functions with Applications to Means,* Journal of Function Spaces and Appl., Volume 2012, Article ID 980438, 14 p., doi:10.1155/2012/980438.

[20] G.S. Yang, D.Y. Hwang, K.L. Tseng, *Some inequalities for differentiable convex and concave mappings,* Comput. Math. Appl. 47 (2004) 207-216.

Kilis 7 Aralik University, Department of Mathematics, 79000, Kilis, TURKEY
E-mail address: `mevluttunc@kilis.edu.tr`